



10-mg Warfarin Initiation Nomogram

Note: The American College of Chest Physicians (ACCP) recommends initiation of warfarin 5mg daily or less in patients who are: elderly, debilitated, malnourished, have heart failure, liver disease, recent major surgery, or taking medications known to increase sensitivity to warfarin (e.g. amiodarone).

Reference: Chest 2008; 133; 160-198.

Day 3 INR	Warfarin Dose on Days 3, 4, mg	Day 5 INR	Warfarin Dose on Days 5, 6, 7, mg
< 1.3	15, 15	< 2.0	15, 15, 15
1.3 – 1.4	10, 10	2.0 – 3.0	7.5, 5, 7.5
		3.1 – 3.5	0, 5, 5
		> 3.5	0, 0, 2.5
1.5 – 1.6	10, 5	< 2.0	7.5, 7.5, 7.5
1.7 – 1.9	5, 5	2.0 – 3.0	5, 5, 5
		3.1 – 3.5	2.5, 2.5, 2.5
		> 3.5	0, 2.5, 2.5
2.0 – 2.2	2.5, 2.5	< 2.0	5, 5, 5
2.3 – 3.0	0, 2.5	2.0 – 3.0	2.5, 5, 2.5
		3.1 – 3.5	0, 2.5, 0
		> 3.5	0, 0, 2.5
> 3.0	0, 0	< 2.0	2.5, 2.5, 2.5
		2.0 – 3.0	2.5, 0, 2.5
		3.1 – 4.0	0, 2.5, 0
		> 4.0	0, 0, 2.5

Reference: Kovacs MJ, et al. Comparison of 10mg and 5mg warfarin initiation nomograms together with low-molecular-weight heparin for outpatient treatment of acute venous thromboembolism.

Annals of Internal Medicine 2003; 138: 714-719.

UNIVERSITY OF UTAH HEALTH CARE THROMBOSIS SERVICE

UNIVERSITY HOSPITAL INPATIENT ACCESS: PAGER: 1- 800- 783 -3735

UNIVERSITY THROMBOSIS CENTER: PHONE: 801- 585-3713; FAX: 801- 585- 7978

REDWOOD ANTICOAGULATION CLINIC: PHONE: 801-213- 9150; FAX 801- 213- 9160